


#9

# APPLICATION UNDER UNITED STATES PATENT LAWS

Atty. Dkt. No. PW 279062  
(M#)

Invention: NEW NUCLEOTIDE SEQUENCES WHICH ENCODE FOR THE citA GENE

Inventor (s): Bettina MÖCKEL  
Mike FARWICK  
Thomas HERMANN  
Achim MARX  
Walter PFEFFERLE



09/804,060

Pillsbury Winthrop LLP  
Intellectual Property Group  
1600 Tysons Boulevard  
McLean, VA 22120  
Attorneys  
Telephone: (703) 905-2173

This is a:

- ☐ Provisional Application  
☒ Regular Utility Application

☐ Continuing Application

☒ The contents of the parent are incorporated  
by reference

☐ PCT National Phase Application

☐ Design Application

☐ Reissue Application

☐ Plant Application

☒ Substitute Specification  
Sub. Spec Filed March 13, 2001

in App. No. 09/804,060

☐ Marked up Specification re  
Sub. Spec. filed  
In App. No. /

## SPECIFICATION

**New nucleotide sequences which code for the citA gene**

The invention provides nucleotide sequences from coryneform bacteria which code for the citA gene and a process for the fermentative preparation of amino acids, in particular L-lysine, by attenuation of the citA gene. The citA gene codes for the sensor kinase Cit A of a two-component system.

Prior art

L-Amino acids, in particular L-lysine, are used in human medicine and in the pharmaceuticals industry, in the foodstuffs industry and very particularly in animal nutrition.

It is known that amino acids are prepared by fermentation from strains of coryneform bacteria, in particular Corynebacterium glutamicum. Because of their great importance, work is constantly being undertaken to improve the preparation processes. Improvements to the process can relate to fermentation measures, such as, for example, stirring and supply of oxygen, or the composition of the nutrient media, such as, for example, the sugar concentration during the fermentation, or the working up to the product form by, for example, ion exchange chromatography, or the intrinsic output properties of the microorganism itself.

Methods of mutagenesis, selection and mutant selection are used to improve the output properties of these microorganisms. Strains which are resistant to antimetabolites or are auxotrophic for metabolites of regulatory importance and which produce amino acids are obtained in this manner.

09304050-075501

Methods of the recombinant DNA technique have also been employed for some years for improving the strain of *Corynebacterium* strains which produce L-amino acids.

#### Object of the invention

- 5 The inventors had the object of providing new measures for improved fermentative preparation of amino acids, in particular L-lysine.

#### Description of the invention

- If L-lysine or lysine are mentioned in the following, this  
10 also means the salts, such as e. g. lysine monohydrochloride or lysine sulfate.

The invention provides an isolated polynucleotide from coryneform bacteria, comprising a polynucleotide sequence which codes for the *citA* gene chosen from the group  
15 consisting of

- a) polynucleotide which is identical to the extent of at least 70 % to a polynucleotide which codes for a polypeptide which comprises the amino acid sequence of SEQ ID No. 2,
- 20 b) polynucleotide which codes for a polypeptide which comprises an amino acid sequence which is identical to the extent of at least 70 % to the amino acid sequence of SEQ ID No. 2,
- c) polynucleotide which is complementary to the  
25 polynucleotides of a) or b), and
- d) polynucleotide comprising at least 15 successive nucleotides of the polynucleotide sequence of a), b) or c),

the polypeptide preferably having the activity of sensor  
30 kinase *CitA*.

00004060 07501  
T05220 05040860

The invention also provides the abovementioned polynucleotide, this preferably being a DNA which is capable of replication, comprising:

- (i) the nucleotide sequence shown in SEQ ID No.1 or
- 5 (ii) at least one sequence which corresponds to sequence (i) within the range of the degeneration of the genetic code, or
- (iii) at least one sequence which hybridizes with the sequences complementary to sequences (i) or (ii),  
10 and optionally
- (iv) sense mutations of neutral function in (i).

The invention also provides:

a polynucleotide, in particular DNA, which is capable of replication and comprises the nucleotide sequence as  
15 shown in SEQ ID No.1;

a polynucleotide which codes for a polypeptide which comprises the amino acid sequence as shown in SEQ ID No. 2;

a vector containing parts of the polynucleotide according  
20 to the invention, but at least 15 successive nucleotides of the sequence claimed

and coryneform bacteria in which the *citA* gene is attenuated, in particular by an insertion or deletion.

The invention also provides polynucleotides which  
25 substantially comprise a polynucleotide sequence, which are obtainable by screening by means of hybridization of a corresponding gene library of a coryneform bacterium, which comprises the complete gene or parts thereof, with a probe which comprises the sequence of the polynucleotide

09804060.072501

according to the invention or a fragment thereof, and isolation of the polynucleotide sequence mentioned.

Polynucleotides comprising the sequences according to the invention are suitable as hybridization probes for RNA,  
5 cDNA and DNA, in order to isolate, in the full length, nucleic acids or polynucleotides or genes which code for CitA protein or to isolate those nucleic acids or polynucleotides or genes which have a high similarity with the sequence of the citA gene.

10 Polynucleotides comprising the sequences according to the invention are furthermore suitable as primers with the aid of which DNA of genes which code for the CitA protein can be prepared by the polymerase chain reaction (PCR).

Such oligonucleotides which serve as probes or primers  
15 comprise at least 30, preferably at least 20, very particularly preferably at least 15 successive nucleotides. Oligonucleotides which have a length of at least 40 or 50 nucleotides are also suitable.

"Isolated" means separated out of its natural environment.

20 "Polynucleotide" in general relates to polyribonucleotides and polydeoxyribonucleotides, it being possible for these to be non-modified RNA or DNA or modified RNA or DNA.

The polynucleotides according to the invention include a polynucleotide according to SEQ ID No. 1 or a fragment  
25 prepared therefrom and also those which are at least 70 %, preferably at least 80 % and in particular at least 90 % to 95 % identical to the polynucleotide according to SEQ ID No. 1 or a fragment prepared therefrom.

"Polypeptides" are understood as meaning peptides or  
30 proteins which comprise two or more amino acids bonded via peptide bonds.

00004060.03040860

The polypeptides according to the invention include a polypeptide according to SEQ ID No. 2, in particular those with the biological activity of the CitA protein and also those which are at least 70 %, preferably at least 80 % and  
5 in particular at least 90 % to 95 % identical to the polypeptide according to SEQ ID No. 2 and have the activity mentioned.

The invention moreover relates to a process for the fermentative preparation of amino acids, in particular L-  
10 lysine, using coryneform bacteria which in particular already produce amino acids and in which the nucleotide sequences which code for the citA gene are attenuated, in particular eliminated or expressed at a low level.

The term "attenuation" in this connection describes the  
15 reduction or elimination of the intracellular activity of one or more enzymes (proteins) in a microorganism which are coded by the corresponding DNA, for example by using a weak promoter or using a gene or allele which codes for a corresponding enzyme with a low activity or inactivates the  
20 corresponding gene or enzyme (protein), and optionally combining these measures.

The microorganisms which the present invention provides can prepare amino acids, in particular L-lysine, from glucose, sucrose, lactose, fructose, maltose, molasses, starch,  
25 cellulose or from glycerol and ethanol. They can be representatives of coryneform bacteria, in particular of the genus Corynebacterium. Of the genus Corynebacterium, there may be mentioned in particular the species Corynebacterium glutamicum, which is known among experts  
30 for its ability to produce L-amino acids.

Suitable strains of the genus Corynebacterium, in particular of the species Corynebacterium glutamicum (C. glutamicum), are in particular the known wild-type strains

09804060 072501

- Corynebacterium glutamicum ATCC13032  
Corynebacterium acetoglutamicum ATCC15806  
Corynebacterium acetoacidophilum ATCC13870  
Corynebacterium melassecola ATCC17965  
5 Corynebacterium thermoaminogenes FERM BP-1539  
Brevibacterium flavum ATCC14067  
Brevibacterium lactofermentum ATCC13869 and  
Brevibacterium divaricatum ATCC14020

or L-amino acid-producing mutants or strains prepared  
10 therefrom, such as, for example, the L-lysine-producing  
strains

- Corynebacterium glutamicum FERM-P 1709  
Brevibacterium flavum FERM-P 1708  
Brevibacterium lactofermentum FERM-P 1712  
15 Corynebacterium glutamicum FERM-P 6463  
Corynebacterium glutamicum FERM-P 6464  
Corynebacterium glutamicum DM58-1  
Corynebacterium glutamicum DG52-5  
Corynebacterium glutamicum DSM 5715 and  
20 Corynebacterium glutamicum DSM 12866

The inventors have succeeded in isolating the new *citA* gene  
of *C. glutamicum* which codes for the Cita protein and which  
is a sensor kinase of a two-component system.

- To isolate the *citA* gene or also other genes of *C.*  
25 *glutamicum*, a gene library of this microorganism is first  
set up in *Escherichia coli* (*E. coli*). The setting up of  
gene libraries is described in generally known textbooks  
and handbooks. The textbook by Winnacker: *Gene und Klone, Eine Einführung in die Gentechnologie* [Genes and Clones, An  
30 Introduction to Genetic Engineering] (Verlag Chemie,  
Weinheim, Germany, 1990), or the handbook by Sambrook et  
al.: *Molecular Cloning, A Laboratory Manual* (Cold Spring  
Harbor Laboratory Press, 1989) may be mentioned as an  
example. A well-known gene library is that of the *E. coli*

K-12 strain W3110 set up in  $\lambda$ -vectors by Kohara et al. (Cell 50, 495-508 (1987)). Bathe et al. (Molecular and General Genetics, 252:255-265, 1996) describe a gene library of *C. glutamicum* ATCC13032, which was set up with the aid of the cosmid vector SuperCos I (Wahl et al., 1987, Proceedings of the National Academy of Sciences USA, 84:2160-2164) in the *E. coli* K-12 strain NM554 (Raleigh et al., 1988, Nucleic Acids Research 16:1563-1575). Börmann et al. (Molecular Microbiology 6(3), 317-326 (1992)) in turn describe a gene library of *C. glutamicum* ATCC13032 using the cosmid pH79 (Hohn and Collins, 1980, Gene 11, 291-298).

To prepare a gene library of *C. glutamicum* in *E. coli* it is also possible to use plasmids such as pBR322 (Bolivar, 1979, Life Sciences, 25, 807-818) or pUC9 (Vieira et al., 1982, Gene, 19:259-268). Suitable hosts are, in particular, those *E. coli* strains which are restriction- and recombination-defective, such as, for example, the strain DH5 $\alpha$  (Jeffrey H. Miller: "A Short Course in Bacterial Genetics, A Laboratory Manual and Handbook for Escherichia coli and Related Bacteria", Cold Spring Harbour [sic] Laboratory Press, 1992).

The long DNA fragments cloned with the aid of cosmids or other  $\lambda$ -vectors can then be subcloned in turn into the usual vectors suitable for DNA sequencing.

Methods of DNA sequencing are described, inter alia, by Sanger et al. (Proceedings of the National Academy of Sciences of the United States of America USA, 74:5463-5467, 1977).

The resulting DNA sequences can then be investigated with known algorithms or sequence analysis programs, such as e.g. that of Staden (Nucleic Acids Research 14, 217-232(1986)), that of Marck (Nucleic Acids Research 16,

T05220-09040860



1829-1836 (1988)) or the GCG program of Butler (Methods of Biochemical Analysis 39, 74-97 (1998)).

The new DNA sequence of *C. glutamicum* which codes for the *citA* gene and which, as SEQ ID No. 1, is a constituent of the present invention has been found in this way. The amino acid sequence of the corresponding protein has furthermore been derived from the present DNA sequence by the methods described above. The resulting amino acid sequence of the *citA* gene product is shown in SEQ ID No. 2.

10 Coding DNA sequences which result from SEQ ID No. 1 by the degeneracy of the genetic code are also a constituent of the invention. In the same way, DNA sequences which hybridize with SEQ ID No. 1 or parts of SEQ ID No. 1 are a constituent of the invention. Conservative amino acid  
15 exchanges, such as e.g. exchange of glycine for alanine or of aspartic acid for glutamic acid in proteins, are furthermore known among experts as "sense mutations" which do not lead to a fundamental change in the activity of the protein, i.e. are of neutral function. It is furthermore  
20 known that changes on the N and/or C terminus of a protein cannot substantially impair or can even stabilize the function thereof. Information in this context can be found by the expert, inter alia, in Ben-Bassat et al. (Journal of Bacteriology 169:751-757 (1987)), in O'Regan et al. (Gene  
25 77:237-251 (1989)), in Sahin-Toth et al. (Protein Sciences 3:240-247 (1994)), in Hochuli et al. (Bio/Technology 6:1321-1325 (1988)) and in known textbooks of genetics and molecular biology. Amino acid sequences which result in a corresponding manner from SEQ ID No. 2 are also a  
30 constituent of the invention.

Finally, DNA sequences which are prepared by the polymerase chain reaction (PCR) using primers which result from SEQ ID No. 1 are a constituent of the invention. Such oligonucleotides typically have a length of at least 15  
35 nucleotides.

09804060.072501

Instructions for identifying DNA sequences by means of hybridization can be found by the expert, inter alia, in the handbook "The DIG System Users Guide for Filter Hybridization" from Boehringer Mannheim GmbH (Mannheim, Germany, 1993) and in Liebl et al. (International Journal of Systematic Bacteriology 41: 255-260 (1991)). The hybridization takes place under stringent conditions, that is to say only hybrids in which the probe and target sequence, i. e. the polynucleotides treated with the probe, are at least 70 % identical are formed. It is known that the stringency of the hybridization, including the washing steps, is influenced or determined by varying the buffer composition, the temperature and the salt concentration. The hybridization reaction is preferably carried out under a relatively low stringency compared with the washing steps (Hybaid Hybridisation Guide, Hybaid Limited, Teddington, UK, 1996). A 5x SSC buffer at a temperature of approx. 50 - 68°C, for example, can be employed for the hybridization reaction. Probes can also hybridize here with polynucleotides which are less than 70 % identical to the sequence of the probe. Such hybrids are less stable and are removed by washing under stringent conditions. This can be achieved, for example, by lowering the salt concentration to 2x SSC and optionally subsequently 0.5x SSC (The DIG System User's Guide for Filter Hybridisation, Boehringer Mannheim, Mannheim, Germany, 1995) a temperature of approx. 50 - 68°C being established. Polynucleotide fragments which are, for example, at least 70 % or at least 80 % or at least 90 % to 95 % identical to the sequence of the probe employed can be isolated by increasing the hybridization temperature stepwise from 50 to 68°C in steps of approx. 1 - 2°C. Further instructions on hybridization are obtainable on the market in the form of so-called kits (e.g. DIG Easy Hyb from Roche Diagnostics GmbH, Mannheim, Germany, Catalogue No. 1603558).

OligonukleotidesA: [sic] a Practical Approach (IRL Press, Oxford, UK, 1984) and in Newton and Graham: PCR (Spektrum Akademischer Verlag, Heidelberg, Germany, 1994).

10 citA gene.

optionally be combined.

20 ribosome binding sites, the start codon and terminators.

(Nucleic Acids Research 26: 3548 (1998), in Jensen and

e.g. the textbook by Knippers ("Molekulare Genetik  
30 [Molecular Genetics]", 6th edition, Georg Thieme Verlag,

Stuttgart, Germany, 1995) or that by Winnacker ("Gene und Klone [Genes and Clones]", VCH Verlagsgesellschaft, Weinheim, Germany, 1990).

- Mutations which lead to a change or reduction in the catalytic properties of enzyme proteins are known from the prior art; examples which may be mentioned are the works by Qiu and Goodman (Journal of Biological Chemistry 272: 8611-8617 (1997)), Sugimoto et al. (Bioscience Biotechnology and Biochemistry 61: 1760-1762 (1997)) and Möckel ("Die Threonindehydratase aus Corynebacterium glutamicum: Aufhebung der allosterischen Regulation und Struktur des Enzyms [Threonine dehydratase from Corynebacterium glutamicum: Cancelling the allosteric regulation and structure of the enzyme]", Reports from the Jülich Research Centre, Jül-2906, ISSN09442952, Jülich, Germany, 1994). Summarizing descriptions can be found in known textbooks of genetics and molecular biology, such as e.g. that by Hagemann ("Allgemeine Genetik [General Genetics]", Gustav Fischer Verlag, Stuttgart, 1986).

- Possible mutations are transitions, transversions, insertions and deletions. Depending on the effect of the amino acid exchange on the enzyme activity, missense mutations or nonsense mutations are referred to. Insertions or deletions of at least one base pair (bp) in a gene lead to frame shift mutations, as a consequence of which incorrect amino acids are incorporated or translation is interrupted prematurely. Deletions of several codons typically lead to a complete loss of the enzyme activity. Instructions on generation of such mutations are prior art and can be found in known textbooks of genetics and molecular biology, such as e.g. the textbook by Knippers ("Molekulare Genetik [Molecular Genetics]", 6th edition, Georg Thieme Verlag, Stuttgart, Germany, 1995), that by Winnacker ("Gene und Klone [Genes and Clones]", VCH Verlagsgesellschaft, Weinheim, Germany, 1990) or that by Hagemann ("Allgemeine Genetik [General Genetics]", Gustav Fischer Verlag, Stuttgart, 1986).

09804060-073501

A common method of mutating genes of *C. glutamicum* is the method of gene disruption and gene replacement described by Schwarzer and Pühler (Bio/Technology 9, 84-87 (1991)).

In the method of gene disruption a central part of the coding region of the gene of interest is cloned in a plasmid vector which can replicate in a host (typically *E. coli*), but not in *C. glutamicum*. Possible vectors are, for example, pSUP301 (Simon et al., Bio/Technology 1, 784-791 (1983)), pK18mob or pK19mob (Schäfer et al., Gene 145, 69-73 (1994)), pK18mobsacB or pK19mobsacB (Jäger et al., Journal of Bacteriology 174: 5462-65 (1992)), pGEM-T (Promega corporation, Madison, WI, USA), pCR2.1-TOPO (Shuman (1994). Journal of Biological Chemistry 269:32678-84; US Patent 5,487,993), pCR®Blunt (Invitrogen, Groningen, Holland; Bernard et al., Journal of Molecular Biology, 234: 534-541 (1993)) or pEM1 (Schrumpf et al, 1991, Journal of Bacteriology 173:4510-4516). The plasmid vector which contains the central part of the coding region of the gene is then transferred into the desired strain of *C. glutamicum* by conjugation or transformation. The method of conjugation is described, for example, by Schäfer et al. (Applied and Environmental Microbiology 60, 756-759 (1994)). Methods for transformation are described, for example, by Thierbach et al. (Applied Microbiology and Biotechnology 29, 356-362 (1988)), Dunican and Shivnan (Bio/Technology 7, 1067-1070 (1989)) and Tauch et al. (FEMS Microbiological Letters 123, 343-347 (1994)). After homologous recombination by means of a "cross-over" event, the coding region of the gene in question is interrupted by the vector sequence and two incomplete alleles are obtained, one lacking the 3' end and one lacking the 5' end. This method has been used, for example, by Fitzpatrick et al. (Applied Microbiology and Biotechnology 42, 575-580 (1994)) to eliminate the *recA* gene of *C. glutamicum*.

- In the method of gene replacement, a mutation, such as e.g. a deletion, insertion or base exchange, is established in vitro in the gene of interest. The allele prepared is in turn cloned in a vector which is not replicative for *C. glutamicum* and this is then transferred into the desired host of *C. glutamicum* by transformation or conjugation. After homologous recombination by means of a first "cross-over" event which effects integration and a suitable second "cross-over" event which effects excision in the target gene or in the target sequence, the incorporation of the mutation or of the allele is achieved. This method was used, for example, by Peters-Wendisch et al. (Microbiology 144, 915 - 927 (1998)) to eliminate the *pyc* gene of *C. glutamicum* by a deletion.
- 15 A deletion, insertion or a base exchange can be incorporated into the *citA* gene in this manner.

- In addition, it may be advantageous for the production of L-amino acids, in particular L-lysine, to enhance, in particular to over-express, one or more enzymes of the particular biosynthesis pathway, of glycolysis, of anaplerosis, of the pentose phosphate cycle or of amino acid export and optionally regulatory proteins, in addition to attenuation of the *citA* gene.

- The term "enhancement" in this connection describes the increase in the intracellular activity of one or more enzymes (proteins) in a microorganism which are coded by the corresponding DNA, for example by increasing the number of copies of the gene or genes, using a potent promoter or using a gene or allele which codes for a corresponding enzyme (protein) having a high activity, and optionally combining these measures.

Thus, for example, for the preparation of L-lysine, at the same time one or more of the genes chosen from the group consisting of

- 105220-09040360
- the dapA gene which codes for dihydrodipicolinate synthase (EP-B 0 197 335),
  - the gap gene which codes for glyceraldehyde 3-phosphate dehydrogenase (Eikmanns (1992), Journal of Bacteriology 5 174:6076-6086),
  - the zwf gene which codes for glucose 6-phosphate dehydrogenase (JP-A-09224661),
  - the pyc gene which codes for pyruvate carboxylase (DE-A-198 31 609),
  - 10 • the lysE gene which codes for lysine export (DE-A-195 48 222)
  - the lysC gene which codes for a feed back resistant aspartate kinase (EP-B-0387527; EP-A-0699759; WO 00/63388), or
  - 15 • the zwal gene which codes for the Zwal protein (DE: 199 59 328.0, DSM 13115)

can be enhanced, in particular over-expressed.

It may furthermore be advantageous for the production of amino acids, in particular L-lysine, in addition to the 20 attenuation of the citA gene, at the same time for one or more of the genes chosen from the group consisting of

- the pck gene which codes for phosphoenol pyruvate carboxykinase (DE 199 50 409.1, DSM 13047),
- the pgi gene which codes for glucose 6-phosphate 25 isomerase (US 09/396,478, DSM 12969),
- the poxB gene which codes for pyruvate oxidase (DE:1995 1975.7, DSM 13114)

- the *zwa2* gene which codes for the Zwa2 protein  
(DE: 199 59,327.2, DSM 13113)

to be attenuated.

In addition to attenuation of the *citA* gene it may  
5 furthermore be advantageous, for the production of amino  
acids, in particular L-lysine, to eliminate undesirable  
side reactions, (Nakayama: "Breeding of Amino Acid  
Producing Microorganisms", in: Overproduction of Microbial  
Products, Krumphanzl, Sikyta, Vanek (eds.), Academic Press,  
10 London, UK, 1982).

The invention also provides the microorganisms prepared  
according to the invention, and these can be cultured  
continuously or discontinuously in the batch process (batch  
culture) or in the fed batch (feed process) or repeated fed  
15 batch process (repetitive feed process) for the purpose of  
production of L-amino acids, in particular L-lysine. A  
summary of known culture methods is described in the  
textbook by Chmiel (Bioprozesstechnik 1. Einführung in die  
Bioverfahrenstechnik [Bioprocess Technology 1. Introduction  
20 to Bioprocess Technology (Gustav Fischer Verlag, Stuttgart,  
1991)) or in the textbook by Storhas (Bioreaktoren und  
periphere Einrichtungen [Bioreactors and Peripheral  
Equipment] (Vieweg Verlag, Braunschweig/Wiesbaden, 1994)).

The culture medium to be used must meet the requirements of  
25 the particular strains in a suitable manner. Descriptions  
of culture media for various microorganisms are contained  
in the handbook "Manual of Methods for General  
Bacteriology" of the American Society for Bacteriology  
(Washington D.C., USA, 1981). Sugars and carbohydrates,  
30 such as e.g. glucose, sucrose, lactose, fructose, maltose,  
molasses, starch and cellulose, oils and fats, such as, for  
example, soya oil, sunflower oil, groundnut oil and coconut  
fat, fatty acids, such as, for example, palmitic acid,  
stearic acid and linoleic acid, alcohols, such as, for

09804050 072501



example, glycerol and ethanol, and organic acids, such as, for example, acetic acid, can be used as the source of carbon. These substances can be used individually or as a mixture.

- 5 Organic nitrogen-containing compounds, such as peptones, yeast extract, meat extract, malt extract, corn steep liquor, soya bean flour and urea, or inorganic compounds, such as ammonium sulfate, ammonium chloride, ammonium phosphate, ammonium carbonate and ammonium nitrate, can be  
10 used as the source of nitrogen. The sources of nitrogen can be used individually or as a mixture.

Phosphoric acid, potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding sodium-containing salts can be used as the source of phosphorus.

- 15 The culture medium must furthermore comprise salts of metals, such as, for example, magnesium sulfate or iron sulfate, which are necessary for growth. Finally, essential growth substances, such as amino acids and vitamins, can be employed in addition to the abovementioned substances.  
20 Suitable precursors can moreover be added to the culture medium. The starting substances mentioned can be added to the culture in the form of a single batch, or can be fed in during the culture in a suitable manner.

- Basic compounds, such as sodium hydroxide, potassium  
25 hydroxide, ammonia or aqueous ammonia, or acid compounds, such as phosphoric acid or sulfuric acid, can be employed in a suitable manner to control the pH of the culture. Antifoams, such as, for example, fatty acid polyglycol esters, can be employed to control the development of foam.  
30 Suitable substances having a selective action, such as, for example, antibiotics, can be added to the medium to maintain the stability of plasmids. To maintain aerobic conditions, oxygen or oxygen-containing gas mixtures, such as, for example, air, are introduced into the culture. The  
35 temperature of the culture is usually 20°C to 45°C, and

00004060-02501

preferably 25°C to 40°C. Culturing is continued until a maximum of the desired product has formed. This target is usually reached within 10 hours to 160 hours.

Methods for the determination of L-amino acids are known from the prior art. The analysis can thus be carried out, for example, as described by Spackman et al. (Analytical Chemistry, 30, (1958), 1190) by anion exchange chromatography with subsequent ninhydrin derivatization, or it can be carried out by reversed phase HPLC, for example as described by Lindroth et al. (Analytical Chemistry (1979) 51: 1167-1174).

The invention furthermore relates to a process for the fermentative preparation of an amino acid chosen from the group consisting of L-asparagine, L-threonine, L-serine, L-glutamate, L-glycine, L-alanine, L-cysteine, L-valine, L-methionine, L-isoleucine, L-leucine, L-tyrosine, L-phenylalanine, L-histidine, L-lysine, L-tryptophan and L-arginine, in particular L-lysine, using coryneform bacteria which in particular already produce one or more of the amino acids mentioned.

*Sull  
A'* The following microorganism was deposited on 23.01.2001 as a pure culture at the Deutsche Sammlung für Mikroorganismen [sic] und Zellkulturen (DSMZ = German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany) in accordance with the Budapest Treaty:

- Escherichia coli strain Top10/pCR2.1citAint as DSM 13998.

The present invention is explained in more detail in the following with the aid of embodiment examples.

The isolation of plasmid DNA from Escherichia coli and all techniques of restriction, Klenow and alkaline phosphatase treatment were carried out by the method of Sambrook et al. (Molecular Cloning. A Laboratory Manual, 1989, Cold Spring

00004060-0750

Harbour [sic] Laboratory Press, Cold Spring Harbor, NY, USA). Methods for transformation of *Escherichia coli* are also described in this handbook.

The composition of the usual nutrient media, such as LB or 5 TY medium, can also be found in the handbook by Sambrook et al.

09604060-072501

ExamplesExample 1

Preparation of a genomic cosmid gene library from *C. glutamicum* ATCC 13032

- 5 Chromosomal DNA from *C. glutamicum* ATCC 13032 was isolated as described by Tauch et al. (1995, Plasmid 33:168-179) and partly cleaved with the restriction enzyme Sau3AI (Amersham Pharmacia, Freiburg, Germany, Product Description Sau3AI, Code no. 27-0913-02). The DNA fragments were
- 10 dephosphorylated with shrimp alkaline phosphatase (Roche Molecular Biochemicals, Mannheim, Germany, Product Description SAP, Code no. 1758250). The DNA of the cosmid vector SuperCos1 (Wahl et al. (1987), Proceedings of the National Academy of Sciences, USA 84:2160-2164), obtained
- 15 from Stratagene (La Jolla, USA, Product Description SuperCos1 Cosmid Vektor Kit, Code no. 251301) was cleaved with the restriction enzyme XbaI (Amersham Pharmacia, Freiburg, Germany, Product Description XbaI, Code no. 27-0948-02) and likewise dephosphorylated with shrimp alkaline
- 20 phosphatase.

- The cosmid DNA was then cleaved with the restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, Product Description BamHI, Code no. 27-0868-04). The cosmid DNA treated in this manner was mixed with the treated ATCC13032
- 25 DNA and the batch was treated with T4 DNA ligase (Amersham Pharmacia, Freiburg, Germany, Product Description T4-DNA-Ligase, Code no. 27-0870-04). The ligation mixture was then packed in phages with the aid of Gigapack II XL Packing Extract (Stratagene, La Jolla, USA, Product Description
- 30 Gigapack II XL Packing Extract, Code no. 200217).

For infection of the *E. coli* strain NM554 (Raleigh et al. 1988, Nucleic Acid Res. 16:1563-1575) the cells were taken up in 10 mM MgSO<sub>4</sub> and mixed with an aliquot of the phage

T05220 09040860

suspension. The infection and titering of the cosmid library were carried out as described by Sambrook et al. (1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor), the cells being plated out on LB agar (Lennox, 5 1955, Virology, 1:190) + 100 µg/ml ampicillin. After incubation overnight at 37°C, recombinant individual clones were selected.

### Example 2

#### Isolation and sequencing of the citA gene

- 10 The cosmid DNA of an individual colony was isolated with the Qiaprep Spin Miniprep Kit (Product No. 27106, Qiagen, Hilden, Germany) in accordance with the manufacturer's instructions and partly cleaved with the restriction enzyme Sau3AI (Amersham Pharmacia, Freiburg, Germany, Product Description Sau3AI, Product No. 27-0913-02). The DNA 15 fragments were dephosphorylated with shrimp alkaline phosphatase (Roche Molecular Biochemicals, Mannheim, Germany, Product Description SAP, Product No. 1758250). After separation by gel electrophoresis, the cosmid 20 fragments in the size range of 1500 to 2000 bp were isolated with the QiaExII Gel Extraction Kit (Product No. 20021, Qiagen, Hilden, Germany).

- The DNA of the sequencing vector pZero-1, obtained from Invitrogen (Groningen, The Netherlands, Product Description 25 Zero Background Cloning Kit, Product No. K2500-01) was cleaved with the restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, Product Description BamHI, Product No. 27-0868-04). The ligation of the cosmid fragments in the sequencing vector pZero-1 was carried out 30 as described by Sambrook et al. (1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor), the DNA mixture being incubated overnight with T4 ligase (Pharmacia Biotech, Freiburg, Germany). This ligation mixture was then electroporated (Tauch et al. 1994, FEMS Microbiol Letters,

000169 00040860

123:343-7) into the E. coli strain DH5 $\alpha$ MCR (Grant, 1990, Proceedings of the National Academy of Sciences, U.S.A., 87:4645-4649) and plated out on LB agar (Lennox, 1955, Virology, 1:190) with 50  $\mu$ g/ml zeocin.

- 5 The plasmid preparation of the recombinant clones was carried out with Biorobot 9600 (Product No. 900200, Qiagen, Hilden, Germany). The sequencing was carried out by the dideoxy chain termination method of Sanger et al. (1977, Proceedings of the National Academies of Sciences, U.S.A., 74:5463-5467) with modifications according to Zimmermann et al. (1990, Nucleic Acids Research, 18:1067). The "RR dRhodamin Terminator Cycle Sequencing Kit" from PE Applied Biosystems (Product No. 403044, Weiterstadt, Germany) was used. The separation by gel electrophoresis and analysis of the sequencing reaction were carried out in a "Rotiphoresis NF Acrylamide/Bisacrylamide" Gel (29:1) (Product No. A124.1, Roth, Karlsruhe, Germany) with the "ABI Prism 377" sequencer from PE Applied Biosystems (Weiterstadt, Germany).
- 20 The raw sequence data obtained were then processed using the Staden program package (1986, Nucleic Acids Research, 14:217-231) version 97-0. The individual sequences of the pZerol derivatives were assembled to a continuous contig. The computer-assisted coding region analysis [sic] were prepared with the XNIP program (Staden, 1986, Nucleic Acids Research, 14:217-231). Further analyses were carried out with the "BLAST search program" (Altschul et al., 1997, Nucleic Acids Research, 25:3389-3402) against the non-redundant databank of the "National Center for Biotechnology Information" (NCBI, Bethesda, MD, USA).

The resulting nucleotide sequence is shown in SEQ ID No. 1. Analysis of the nucleotide sequence showed an open reading frame of 1653 bp, which was called the citA gene. The citA gene codes for a polypeptide of of [sic] 551 amino acids.

Example 3

Preparation of an integration vector for integration mutagenesis of the citA gene

From the strain ATCC 13032, chromosomal DNA was isolated by  
5 the method of Eikmanns et al. (Microbiology 140: 1817 -  
1828 (1994)). On the basis of the sequence of the citA gene  
known for *C. glutamicum* from example 2, the following  
oligonucleotides were chosen for the polymerase chain  
reaction (see SEQ ID No. 4 and SEQ ID No. 5):

10 citA-int1:

5' TTC CAG TCG GTG AGG TCA GT 3'

citA-int2:

5' GTA CGA TCG CGG ATG GTT AC 3'

The primers shown were synthesized by MWG Biotech  
15 (Ebersberg, Germany) and the PCR reaction was carried out  
by the standard PCR method of Innis et al. (PCR protocols.  
A guide to methods and applications, 1990, Academic Press)  
with the Taq-polymerase from Boehringer Mannheim (Germany,  
Product Description Taq DNA polymerase, Product No. 1 146  
20 165). With the aid of the polymerase chain reaction, the  
primers allow amplification of an internal fragment of the  
citA gene 480 bp in size. The product amplified in this way  
was tested electrophoretically in a 0.8% agarose gel.

The amplified DNA fragment (see SEQ ID No. 3) was ligated  
25 with the TOPO TA Cloning Kit from Invitrogen Corporation  
(Carlsbad, CA, USA; Catalogue Number K4500-01) in the  
vector pCR2.1-TOPO (Mead et al. (1991) Bio/Technology  
9:657-663).

The *E. coli* strain TOP10 was then electroporated with the  
30 ligation batch (Hanahan, In: DNA cloning. A practical  
approach. Vol.I. IRL-Press, Oxford, Washington DC, USA,  
1985). Selection for plasmid-carrying cells was made by  
plating out the transformation batch on LB agar (Sambrook

09804060-072501

et al., Molecular cloning: a laboratory manual. 2<sup>nd</sup> Ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989), which had been supplemented with 50 mg/l kanamycin. Plasmid DNA was isolated from a transformant 5 with the aid of the QIAprep Spin Miniprep Kit from Qiagen and checked by restriction with the restriction enzyme EcoRI and subsequent agarose gel electrophoresis (0.8%). The plasmid was called pCR2.1citAint and is shown in figure 1.

Sub  
A2  
10 The following microorganism was deposited as a pure culture on 23.01.2001 at the Deutsche Sammlung für Mikroorganismen und Zellkulturen (DSMZ = German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany) in accordance with the Budapest Treaty:

15 • Escherichia coli Top10/pCR2.1citAint as DSM 13998

#### Example 4

Integration mutagenesis of the citA gene in the strains DSM 5715 and FERM-BP 1763

The vector pCR2.1citAint mentioned in example 3 was 20 electroporated by the electroporation method of Tauch et al. (FEMS Microbiological Letters, 123:343-347 (1994)) into the strains Corynebacterium glutamicum DSM 5715 and Brevibacterium lactofermentum FERM-BP 1763. The strain DSM 5715 is an AEC-resistant lysine producer (EP-B-0435132), 25 and the strain FERM-BP 1763 is a mycophenolic acid-resistant valine producer (US-A-5,188,948). The vector pCR2.1citAint cannot replicate independently in DSM5715 and FERM-BP 1763 and is retained in the cell only if it has integrated into the chromosome of DSM 5715 or FERM-BP 1763. 30 Selection of clones with pCR2.1citAint integrated into the chromosome was carried out by plating out the electroporation batch on LB agar (Sambrook et al., Molecular cloning: a laboratory manual. 2<sup>nd</sup> Ed. Cold Spring



Harbor Laboratory Press, Cold Spring Harbor, N.Y.), which had been supplemented with 15 mg/l kanamycin.

For detection of the integration, the citAint fragment was labelled with the Dig hybridization kit from Boehringer by the method of "The DIG System Users Guide for Filter Hybridization" of Boehringer Mannheim GmbH (Mannheim, Germany, 1993). Chromosomal DNA of in each case a potential integrant was isolated by the method of Eikmanns et al. (Microbiology 140: 1817 - 1828 (1994)) and in each case cleaved with the restriction enzymes EcoRI, BamHI and HindIII. The fragments formed were separated by means of agarose gel electrophoresis and hybridized at 68°C with the Dig hybridization [sic] kit from Boehringer. The plasmid pCR2.1citAint mentioned in example 3 had been inserted into the chromosome of DSM5715 and into the chromosome of FERM-BP 1763 within the chromosomal citA gene. The strains were called DSM5715::pCR2.1citAint and FERM-BP 1763::pCR2.1citAint.

#### Example 5

#### 20 Preparation of lysine

The C.glutamicum strain DSM5715::pCR2.1citAint obtained in example 4 was cultured in a nutrient medium suitable for the production of lysine and the lysine content in the culture supernatant was determined.

25 For this, the strain was first incubated on an agar plate with the corresponding antibiotic (brain-heart agar with kanamycin (25 mg/l) [sic] for 24 hours at 33°C. Starting from this agar plate culture, a preculture was seeded (10 ml medium in a 100 ml conical flask). The complete  
30 medium CgIII was used as the medium for the preculture.

09804060-072501

## Medium Cg III

NaCl	2.5 g/l
Bacto-Peptone	10 g/l
Bacto-Yeast extract	10 g/l
Glucose (autoclaved separately)	2% (w/v)

The pH is brought to pH 7.4

Kanamycin (25 mg/l) was added to this. The preculture was incubated for 16 hours at 33°C at 240 rpm on a shaking machine. A main culture was seeded from this preculture such that the initial OD (660 nm) of the main culture is 0.1 OD. Medium MM was used for the main culture.

## Medium MM

CSL (corn steep liquor)	5 g/l
MOPS (morpholinopropanesulfonic acid)	20 g/l
Glucose (autoclaved separately)	50g/l
Salts:	
(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub> [sic]	25 g/l
KH <sub>2</sub> PO <sub>4</sub>	0.1 g/l
MgSO <sub>4</sub> * 7 H <sub>2</sub> O	1.0 g/l
CaCl <sub>2</sub> * 2 H <sub>2</sub> O	10 mg/l
FeSO <sub>4</sub> * 7 H <sub>2</sub> O	10 mg/l
MnSO <sub>4</sub> * H <sub>2</sub> O	5.0mg/l
Biotin (sterile-filtered)	0.3 mg/l

T05240"09040860

Thiamine \* HCl (sterile-filtered) 0.2 mg/l  
Leucine (sterile-filtered) 0.1 g/l  
CaCO<sub>3</sub> 25 g/l

The CSL, MOPS and the salt solution were brought to pH 7 with aqueous ammonia and autoclaved. The sterile substrate and vitamin solutions were then added, and the CaCO<sub>3</sub> autoclaved in the dry state was added.

- 5 Culturing was carried out in a 10 ml volume in a 100 ml conical flask with baffles. Kanamycin (25 mg/l) was added. Culturing was carried out at 33°C and 80% atmospheric humidity.

- After 72 hours, the OD was determined at a measurement  
10 wavelength of 660 nm with a Biomek 1000 (Beckmann Instruments GmbH, Munich). The amount of lysine formed was determined with an amino acid analyzer from Eppendorf-BioTronik (Hamburg, Germany) by ion exchange chromatography and post-column derivatization with ninhydrin detection.

- 15 The result of the experiment is shown in table 1.

Table 1

Strain	OD(660)	Lysine HCl g/l
DSM5715	7.5	13.3
DSM5715::pCR2.1citAint	7.4	14.4

Example 6

## Preparation of valine

The *B. lactofermentum* strain FERM-BP 1763::pCR2.1citAint obtained in example 4 was cultured in a nutrient medium suitable for the production of valine and the valine content in the culture supernatant was determined.

For this, the strain was first incubated on an agar plate with the corresponding antibiotic (brain-heart agar with kanamycin (25 mg/l)[sic] for 24 hours at 33°C. Starting from this agar plate culture, a preculture was seeded (10 ml medium in a 100 ml conical flask). The complete medium CgIII was used as the medium for the preculture. Kanamycin (25 mg/l) was added to this. The preculture was incubated for 16 hours at 33°C at 240 rpm on a shaking machine. A main culture was seeded from this preculture such that the initial OD (660nm) of the main culture was 0.1 OD. Medium MM was used for the main culture.

## Medium MM

CSL (corn steep liquor)	5 g/l
MOPS (morpholinopropanesulfonic acid)	20 g/l
Glucose (autoclaved separately)	50g/l
Salts:	
(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub> [sic]	25 g/l
KH <sub>2</sub> PO <sub>4</sub>	0.1 g/l
MgSO <sub>4</sub> * 7 H <sub>2</sub> O	1.0 g/l
CaCl <sub>2</sub> * 2 H <sub>2</sub> O	10 mg/l

09804060-072501

FeSO <sub>4</sub> * 7 H <sub>2</sub> O	10 mg/l
MnSO <sub>4</sub> * H <sub>2</sub> O	5.0mg/l
Isoleucine (sterile-filtered)	0.1 g/l
Methionine (sterile-filtered)	0.1 g/l
Thiamine * HCl (sterile-filtered)	0.2 mg/l
Leucine (sterile-filtered)	0.1 g/l
CaCO <sub>3</sub>	25 g/l

The CSL (corn steep liquor), MOPS (morpholinopropanesulfonic acid) and the salt solution were brought to pH 7 with aqueous ammonia and autoclaved. The sterile substrate and vitamin solutions were then added, 5 and the CaCO<sub>3</sub> autoclaved in the dry state was added.

Culturing was carried out in a 10 ml volume in a 100 ml conical flask with baffles. Kanamycin (25 mg/l) was added. Culturing was carried out at 33°C and 80% atmospheric humidity.

- 10 After 48 hours, the OD was determined at a measurement wavelength of 660 nm with a Biomek 1000 (Beckmann Instruments GmbH, Munich). The amount of valine formed was determined with an amino acid analyzer from Eppendorf-BioTronik (Hamburg, Germany) by ion exchange chromatography 15 and post-column derivatization with ninhydrin detection.

The result of the experiment is shown in table 2.

09804060-072501

Table 2

Strain	OD(660)	Valine HCl g/l
FERM-BP 1763	12.1	7.5
FERM-BP 1763::pCR2.1citAint	13.5	10.8

105270" 03040860

## SEQUENCE PROTOCOL

&lt;110&gt; Degussa-Hüls AG

5 &lt;120&gt; New nucleotide sequences which code for the citA gene

&lt;130&gt; 000169 BT

&lt;140&gt;

10 &lt;141&gt;

&lt;160&gt; 5

&lt;170&gt; Patent In Ver. 2.1

15

&lt;210&gt; 1

&lt;211&gt; 2055

&lt;212&gt; DNA

&lt;213&gt; Corynebacterium glutamicum

20

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (201)..(1853)

&lt;223&gt; citA gene

25

&lt;400&gt; 1

tttctgtgtt tctcgaaact ttgagatccc gagtggtctg tgttgcttgt gggagtataa 60

gggtggcgcggt gtcacgcaca gaagtgtttg gtgcattgcc tgaggtagtg cgcaaaataa 120

30

gacttttctg cattatgatc agaattgttg gcctgggact tgccttcacg ctctgctgat 180

aatcgccccc gggggtagac atg tct gtt ggt gga tcc gac tgg aaa aac ttc 233

Met Ser Val Gly Gly Ser Asp Trp Lys Asn Phe  
1 5 10

35

aag gag gtg gac atc att cgc ttt gct acc cga ata ctg gtg att caa 281

Lys Glu Val Asp Ile Ile Arg Phe Ala Thr Arg Ile Leu Val Ile Gln  
15 20 25

40

gtg gct acc gtc gcg ttg gtg gta gct att tgc acc gga att ttc gca 329

Val Ala Thr Val Ala Leu Val Val Ala Ile Cys Thr Gly Ile Phe Ala  
30 35 40

45

gtt ttg atg atg gat cag atg aaa act gag gcc gag cac aca gcg ctg 377

Val Leu Met Met Asp Gln Met Lys Thr Glu Ala Glu His Thr Ala Leu  
45 50 55

50

tcc atc gga cgt tcg gtg gca tcc aac ccg cag atc cgc gag gaa gta 425

Ser Ile Gly Arg Ser Val Ala Ser Asn Pro Gln Ile Arg Glu Glu Val  
60 65 70 75

55

gcg ctt gat act caa aca gga gca aac cca tcg gcc gaa gaa tta gcc 473

Ala Leu Asp Thr Gln Thr Gly Ala Asn Pro Ser Ala Glu Glu Leu Ala  
80 85 90

gat gga gat atc caa gcg gtt gca cag gcg gcc aat gaa cgc act gga 521

Asp Gly Asp Ile Gln Ala Val Ala Gln Ala Ala Asn Glu Arg Thr Gly  
95 100 105

T05220-09040860

sub A3

5 gct ttg ttt gtc gtt atc act gac ggt tta ggt atc cgc ctg tcc cac 569  
 Ala Leu Phe Val Val Ile Thr Asp Gly Leu Gly Ile Arg Leu Ser His  
 110 115 120  
 10 cca gat gag gaa cgt ctg ggg gag cag gtg agc act agc ttt gag gct 617  
 Pro Asp Glu Glu Arg Leu Gly Glu Gln Val Ser Thr Ser Phe Glu Ala  
 125 130 135  
 15 gcc atg cgg ggt gaa gaa acc atg gcg tgg gag act ggg acc ctc ggt 665  
 Ala Met Arg Gly Glu Glu Thr Met Ala Trp Glu Thr Gly Thr Leu Gly  
 140 145 150 155  
 20 gcg tcc gcg cga gca aaa gtg cct atc ttt gcg ccg gat tct agt gtt 713  
 Ala Ser Ala Arg Ala Lys Val Pro Ile Phe Ala Pro Asp Ser Ser Val  
 160 165 170  
 25 cca gtc ggt gag gtc agt gtt ggg ttt gag cga gac agt gtg tat tcc 761  
 Pro Val Gly Glu Val Ser Val Gly Phe Glu Arg Asp Ser Val Tyr Ser  
 175 180 185  
 30 cgc ctg ccc atg ttc ctc gcc gcc ctt gct ctt att tct gtg ttg gga 809  
 Arg Leu Pro Met Phe Leu Ala Ala Leu Ala Leu Ile Ser Val Leu Gly  
 190 195 200  
 35 atc ctt atc ggc gtg ggt gta gcc atg ggc atg cga cgc cgt tgg gaa 857  
 Ile Leu Ile Gly Val Gly Val Ala Met Gly Met Arg Arg Arg Trp Glu  
 205 210 215  
 40 cgc gtg acc ttg ggt ttg cag ccg gag gag cta gtg acc ctt gtg caa 905  
 Arg Val Thr Leu Gly Leu Gln Pro Glu Glu Leu Val Thr Leu Val Gln  
 220 225 230 235  
 45 aat cag act gca gtc atc gat ggc att gat gag ggc gtg ctg gcg ctg 953  
 Asn Gln Thr Ala Val Ile Asp Gly Ile Asp Glu Gly Val Leu Ala Leu  
 240 245 250  
 50 agc cca aac gga aca att ggg gtg cat aat gag cag gcg caa tcc atg 1001  
 Ser Pro Asn Gly Thr Ile Gly Val His Asn Glu Gln Ala Gln Ser Met  
 255 260 265  
 55 att ggt gca ggt cct atg agt ggc agg acg ttg aaa gaa cta ggg ctt 1049  
 Ile Gly Ala Gly Pro Met Ser Gly Arg Thr Leu Lys Glu Leu Gly Leu  
 270 275 280  
 60 gac ctg ggt ctt gat ggc gtt gta ttg cat ggt cag cat ccg gaa acc 1097  
 Asp Leu Gly Leu Asp Gly Val Val Leu His Gly Gln His Pro Glu Thr  
 285 290 295  
 65 gtt gcc cat aac ggc agg atc ctc tat ctg gat ttc cac ccc gtg cgc 1145  
 Val Ala His Asn Gly Arg Ile Leu Tyr Leu Asp Phe His Pro Val Arg  
 300 305 310 315  
 70 cgt ggg gat caa gat tta ggc tac gtg gta acc atc cgc gat cgt acc 1193  
 Arg Gly Asp Gln Asp Leu Gly Tyr Val Val Thr Ile Arg Asp Arg Thr  
 320 325 330  
 75 gac atc att gaa ctc agt gaa cgc ctc gac tct gtg cgc acc atg acc 1241  
 Asp Ile Ile Glu Leu Ser Glu Arg Leu Asp Ser Val Arg Thr Met Thr

09304060-072501



335 340 345

5 cgc gca ctc cgc gcc cag cgc cac gag ttt gcc aac cgc atc cac acc 1289  
His Ala Leu Arg Ala Gln Arg His Glu Phe Ala Asn Arg Ile His Thr  
350 355 360

10 gca aca ggg ctt atc gac gcc ggc cgc gtc cac gac gcg gca gag ttt 1337  
Ala Thr Gly Leu Ile Asp Ala Gly Arg Val His Asp Ala Ala Glu Phe  
365 370 375

15 cta ggc gat ata tcc cgc aac ggg gga cag tca cat cca ttg atc gga 1385  
Leu Gly Asp Ile Ser Arg Asn Gly Gly Gln Ser His Pro Leu Ile Gly  
380 385 390 395

20 tca gcg cac ctc aat gaa gca ttt ttg agc tca ttt tta agt act gct 1433  
Ser Ala His Leu Asn Glu Ala Phe Leu Ser Ser Phe Leu Ser Thr Ala  
400 405 410

25 tct att tcg gca tct gaa aag ggc gtt agt ctg cgc atc aac tct gac 1481  
Ser Ile Ser Ala Ser Glu Lys Gly Val Ser Leu Arg Ile Asn Ser Asp  
415 420 425

30 acg ctc atc ctt ggc aat gtt aaa gat cca gaa gat gta gca acc att 1529  
Thr Leu Ile Leu Gly Thr Val Lys Asp Pro Glu Asp Val Ala Thr Ile  
430 435 440

35 ttg ggt aat tta atc aac aat gcc atc gac gcc gcg gtg gca ggt gaa 1577  
Leu Gly Asn Leu Ile Asn Asn Ala Ile Asp Ala Val Ala Gly Glu  
445 450 455

40 gcc cca cgg tgg att gag ctt acg ttg atg gat gat gcc gat acg ctg 1625  
Ala Pro Arg Trp Ile Glu Leu Thr Leu Met Asp Asp Ala Asp Thr Leu  
460 465 470 475

45 gtc att tct gtt gca gat tct ggt cct gga atc cca gag ggc gtg gat 1673  
Val Ile Ser Val Ala Asp Ser Gly Pro Gly Ile Pro Glu Gly Val Asp  
480 485 490

50 gta ttt gcc aca gcc acc cag ata gga gac tct gaa gat aat gaa cgc 1721  
Val Phe Ala Thr Ala Thr Gln Ile Gly Asp Ser Glu Asp Asn Glu Arg  
495 500 505

55 acc cac ggg cat ggc att ggt cta aaa ctg tgc cgg gct ttg gct aga 1769  
Thr His Gly His Gly Ile Gly Leu Lys Leu Cys Arg Ala Leu Ala Arg  
510 515 520

60 tca cat ggt ggc gat gtc tgg gtg att gat aga gga acc gaa gat ggc 1817  
Ser His Gly Gly Asp Val Trp Val Ile Asp Arg Gly Thr Glu Asp Gly  
525 530 535

65 gct gta ttt gga gtg aaa cta ccg gga gta atg gag taatgatca 1863  
Ala Val Phe Gly Val Lys Leu Pro Gly Val Met Glu  
540 545 550

70 aacacttaaa gttttagtaa ttgatgatga tttccgcgtc gccggcattc acgcctccat 1923  
cgttgatgcg tcccctggat tttcgggtggt cgggtaccgcg cgtaccctcg cagaggcaaa 1983  
aaccctgata gccacatfff ccccggatct cctacttggt gatgtctacc tccccgacgg 2043

T05220"09040860

cgatggcatt ga

5 <210> 2  
 <211> 551  
 <212> PRT  
 <213> Corynebacterium glutamicum

10 <400> 2  
 Met Ser Val Gly Gly Ser Asp Trp Lys Asn Phe Lys Glu Val Asp Ile  
 1 5 10 15  
 Ile Arg Phe Ala Thr Arg Ile Leu Val Ile Gln Val Ala Thr Val Ala  
 15 20 25 30  
 Leu Val Val Ala Ile Cys Thr Gly Ile Phe Ala Val Leu Met Met Asp  
 35 40 45  
 20 Gln Met Lys Thr Glu Ala Glu His Thr Ala Leu Ser Ile Gly Arg Ser  
 50 55 60  
 Val Ala Ser Asn Pro Gln Ile Arg Glu Glu Val Ala Leu Asp Thr Gln  
 65 70 75 80  
 25 Thr Gly Ala Asn Pro Ser Ala Glu Glu Leu Ala Asp Gly Asp Ile Gln  
 85 90 95  
 Ala Val Ala Gln Ala Ala Asn Glu Arg Thr Gly Ala Leu Phe Val Val  
 30 100 105 110  
 Ile Thr Asp Gly Leu Gly Ile Arg Leu Ser His Pro Asp Glu Glu Arg  
 115 120 125  
 35 Leu Gly Glu Gln Val Ser Thr Ser Phe Glu Ala Ala Met Arg Gly Glu  
 130 135 140  
 Glu Thr Met Ala Trp Glu Thr Gly Thr Leu Gly Ala Ser Ala Arg Ala  
 40 145 150 155 160  
 Lys Val Pro Ile Phe Ala Pro Asp Ser Ser Val Pro Val Gly Glu Val  
 165 170 175  
 45 Ser Val Gly Phe Glu Arg Asp Ser Val Tyr Ser Arg Leu Pro Met Phe  
 180 185 190  
 Leu Ala Ala Leu Ala Leu Ile Ser Val Leu Gly Ile Leu Ile Gly Val  
 195 200 205  
 50 Gly Val Ala Met Gly Met Arg Arg Arg Trp Glu Arg Val Thr Leu Gly  
 210 215 220  
 Leu Gln Pro Glu Glu Leu Val Thr Leu Val Gln Asn Gln Thr Ala Val  
 225 230 235 240  
 55 Ile Asp Gly Ile Asp Glu Gly Val Leu Ala Leu Ser Pro Asn Gly Thr  
 245 250 255  
 Ile Gly Val His Asn Glu Gln Ala Gln Ser Met Ile Gly Ala Gly Pro

0904060.0250

260 265 270  
 Met Ser Gly Arg Thr Leu Lys Glu Leu Gly Leu Asp Leu Gly Leu Asp  
 275 280 285  
 5 Gly Val Val Leu His Gly Gln His Pro Glu Thr Val Ala His Asn Gly  
 290 295 300  
 10 Arg Ile Leu Tyr Leu Asp Phe His Pro Val Arg Arg Gly Asp Gln Asp  
 305 310 315 320  
 Leu Gly Tyr Val Val Thr Ile Arg Asp Arg Thr Asp Ile Ile Glu Leu  
 325 330 335  
 15 Ser Glu Arg Leu Asp Ser Val Arg Thr Met Thr His Ala Leu Arg Ala  
 340 345 350  
 Gln Arg His Glu Phe Ala Asn Arg Ile His Thr Ala Thr Gly Leu Ile  
 355 360 365  
 20 Asp Ala Gly Arg Val His Asp Ala Ala Glu Phe Leu Gly Asp Ile Ser  
 370 375 380  
 25 Arg Asn Gly Gly Gln Ser His Pro Leu Ile Gly Ser Ala His Leu Asn  
 385 390 395 400  
 Glu Ala Phe Leu Ser Ser Phe Leu Ser Thr Ala Ser Ile Ser Ala Ser  
 405 410 415  
 30 Glu Lys Gly Val Ser Leu Arg Ile Asn Ser Asp Thr Leu Ile Leu Gly  
 420 425 430  
 Thr Val Lys Asp Pro Glu Asp Val Ala Thr Ile Leu Gly Asn Leu Ile  
 435 440 445  
 35 Asn Asn Ala Ile Asp Ala Ala Val Ala Gly Glu Ala Pro Arg Trp Ile  
 450 455 460  
 40 Glu Leu Thr Leu Met Asp Asp Ala Asp Thr Leu Val Ile Ser Val Ala  
 465 470 475 480  
 Asp Ser Gly Pro Gly Ile Pro Glu Gly Val Asp Val Phe Ala Thr Ala  
 485 490 495  
 45 Thr Gln Ile Gly Asp Ser Glu Asp Asn Glu Arg Thr His Gly His Gly  
 500 505 510  
 Ile Gly Leu Lys Leu Cys Arg Ala Leu Ala Arg Ser His Gly Gly Asp  
 515 520 525  
 50 Val Trp Val Ile Asp Arg Gly Thr Glu Asp Gly Ala Val Phe Gly Val  
 530 535 540  
 55 Lys Leu Pro Gly Val Met Glu  
 545 550

<211> 481  
 <212> DNA  
 <213> Corynebacterium glutamicum  
 5 <220>  
 <223> citAint  
 <400> 3  
 10 ttccagtcgg tgaggtcagt gttgggtttg agcgagacag tgtgtattcc cgctgcca 60  
 tgttcctcgc cgcccttget cttatttctg tgttgggaat cttatcggc gtgggtgtag 120  
 ccatgggcatt gcgacgccgt tgggaacgcg tgaccttggg tttgcagccg gaggagctag 180  
 tgacccttgt gcaaaatcag actgcagtca tcgatggcat tgatgagggc gtgctggcgc 240  
 tgagcccaaa gggaacaatt ggggtgcata atgagcaggc gcaatccatg attggtgcag 300  
 15 gtcctatgag tggcaggacg ttgaaagaac tagggcttga cctgggtctt gatggcggtt 360  
 tattgcatgg taagcatccg gaaaccgttg ccataacgg caggatcctc tatctggatt 420  
 tccaccccggt gcgccgtggg gatcaagatt taggctacgt ggtaaccatc cgcatcgta 480  
 c 481  
 20 <210> 4  
 <211> 20  
 <212> DNA  
 <213> Corynebacterium glutamicum  
 25 <220>  
 <223> Primer citA-int1  
 <400> 4  
 30 ttccagtcgg tgaggtcagt 20  
 <210> 5  
 <211> 20  
 <212> DNA  
 35 <213> Corynebacterium glutamicum  
 <220>  
 <223> Primer citA-int2  
 40 <400> 5  
 gtacgatcgc ggatgggttac 20

09804050.07501

3630

KK  
12/3/02

The following figure is attached:

Figure 1: Map of the plasmid pCR2.1citAint.

The abbreviations and designations used have the following meaning.

KmR:	Kanamycin resistance gene
EcoRI:	Cleavage site of the restriction enzyme EcoRI
HindIII:	Cleavage site of the restriction enzyme HindIII
BamHI:	Cleavage site of the restriction enzyme BamHI
citAint:	Internal fragment of the citA gene
ColE1:	Replication origin of the plasmid ColE1

5

10

15

T03220"09040860